



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> :  A01N</p>	<p>A2</p>	<p>(11) International Publication Number: <b>WO 99/63813</b>  (43) International Publication Date: 16 December 1999 (16.12.99)</p>
<p>(21) International Application Number: PCT/EP99/03883  (22) International Filing Date: 4 June 1999 (04.06.99)  (30) Priority Data: 9812331.8 8 June 1998 (08.06.98) GB 9903669.1 17 February 1999 (17.02.99) GB  (71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).  (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).  (72) Inventors; and (75) Inventors/Applicants (for US only): ZURFLÜH, René [CH/CH]; Dachslenbergstrasse 54, CH-8180 Bülach (CH). LEADBITTER, Neil [GB/CH]; Therwilerstrasse 117, CH-4104 Oberwil (CH).  (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent &amp; Trademark Dept., CH-4002 Basel (CH).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: FUNGICIDAL COMBINATIONS COMPRISING GLYOXALIC ACID METHYL ESTER-O-METHYLOXIME DERIVATIVES</p> <p>(57) Abstract</p> <p>A method of combating phytopathogenic diseases on crop plants which comprises applying to the crop plants or the locus thereof being infested with said phytopathogenic disease an effective amount of a combination of a) 2-[α-[[[(α-methyl-3-trifluoromethyl-benzyl)imino]-oxy]-o-tolyl]-glyoxalic acid methyl ester-O-methyloxime (I) in association with b) a broad variety of other plant fungicides is particularly effective in combating or preventing diseases of crop plants. These combinations exhibit synergistic fungicidal activity.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

FUNGICIDAL COMBINATIONS COMPRISING GLYOXALIC ACID METHYL ESTER-O-METHYLOXIME DERIVATIVES


The present invention relates to novel fungicidal compositions for the treatment of phytopathogenic diseases of crop plants, especially phytopathogenic fungi, and to a method of combating phytopathogenic diseases on crop plants.

It is known that certain strobilurin derivatives have biological activity against phytopathogenic fungi, e.g. from EP-A-460575 where their properties and methods of preparation are described. On the other hand anilide, carbamate and aminoacid amide fungicides are widely known as plant fungicides for application in various crops of cultivated plants. However, crop tolerance and activity against phytopathogenic plant fungi do not always satisfy the needs of agricultural practice in many incidents and aspects.

It has now been found that the use of


- a) 2-[ $\alpha$ -{[( $\alpha$ -methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acid methyl ester-O-methyloxime, compound I (EP-460575)  
in association with

- b) either a compound of formula IIA




an anilide of formula IIB (EP-545099)

- 2 -



wherein  $R_1$  is fluorine or chlorine; or

a carbamate of formula IIC (WO-96/01256 and WO-96/01258)



wherein X is N or CH, and  $R_2$  is 4-CH<sub>3</sub>, 4-Cl or 2,4-dichloro; or


a compound IID (EP-278595)

methyl(2)-2-{2-[6-(trifluoromethyl)pyrid-2-yloxymethyl]-phenyl}-3-methoxyacrylate; or

a compound IIE (EP-477631)

(E)-N-methyl-2-[2-(2,5-dimethylphenoxy)methyl]phenyl]-2-methoxy-iminoacetamide; or


a compound of formula IIF (WO-95/21154)



wherein  $R_3$  is methyl or ethyl; or


a (S)-valinamide of formula IIG (EP-398072, EP-610764, DE-4321897, WO-96/07638)

- 3 -



wherein  $R_4$  is isopropyl, sec.-butyl or tert.-butyl, and  
 $R_5$  is 4-chlorophenyl, 4-methylphenyl, 4-methoxyphenyl or  $\beta$ -naphthyl, and  
 wherein the asymmetric center is preferably (R); or

a (S)-valinamide of formula IIH (WO-94/25432, WO-96/04252)




wherein  $R_6$  is isopropyl, sec.-butyl or tert.-butyl,  $R_7$  is halogen, methyl or methoxy and  
 $n$  is 0, 1 or 2; or

a compound IIJ (EP-596254)

N-methyl-2-[2-( $\alpha$ -methyl-3-(trifluoromethyl)benzyloximinomethyl)phenyl]-2-methoximinoacetamide; or

a compound of formula IIK (EP-381330)



wherein  $R_8$  is halogen or  $C_1$ - $C_4$ -alkyl, preferably chlorine; or


a compound II L

N-(3'-(1'-chloro-3-methyl 2'-oxopentan))-3,5-dichloro-4-methylbenzamide (EP-600629); or


a compound II M (EP-551048 and WO 96/03044)

(S)-1-anilino-4-methyl-2-methylthio-4-phenylimidazolin-5-one; or

a compound of formula IIN (WO 98/25465)



a compound of formula IIP (WO98/20003)



a compound IIQ

N-methyl-2-[ $\alpha$ -{[( $\alpha$ -methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acidamide-O-methyloxime (EP 569384)

is particularly effective in combating or preventing fungal diseases of crop plants. These combinations exhibit synergistic fungicidal activity.

The combinations according to the invention may also comprise more than one of the active components b) , if broadening of the spectrum of disease control is desired.

The active ingredient combinations are effective against phytopathogenic fungi belonging to the following classes: Ascomycetes (e.g. *Venturia*, *Podosphaera*, *Erysiphe*, *Monilinia*, *Mycosphaerella*, *Uncinula*); Basidiomycetes (e.g. the genus *Hemileia*, *Rhizoctonia*, *Puccinia*); Fungi imperfecti (e.g. *Botrytis*, *Helminthosporium*, *Rhynchosporium*, *Fusarium*, *Septoria*, *Cercospora*, *Alternaria*, *Pycularia* and *Pseudocercospora herpotrichoides*)

(*Tapesia spp.*); Oomycetes (e.g. *Phytophthora*, *Peronospora*, *Bremia*, *Pythium*, *Plasmopara*).

Target crops for the areas of indication disclosed herein comprise within the scope of this invention e.g. the following species of plants: cereals (wheat, barley, rye, oats, rice, sorghum and related crops); beet (sugar beet and fodder beet); pomes, stone fruit and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries); leguminous plants (beans, lentils, peas, soybeans); oil plants (rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans, groundnuts); cucumber plants (marrows, cucumbers, melons); fibre plants (cotton, flax, hemp, jute); citrus fruit (oranges, lemons, grapefruit, mandarins); vegetables (spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, paprika); lauraceae (avocados, cinnamon, camphor); or plants such as maize, tobacco, nuts, coffee, sugar cane, tea, vines, hops, bananas and natural rubber plants, as well as ornamentals (flowers, shrubs, broad-leaved trees and evergreens, such as conifers). This list does not represent any limitation.

The combinations according to the present invention are particularly effective against *Phytophthora*, *Peronospora*, *Bremia*, *Pythium* and *Plasmopara*, in particular against pathogens of monocotyledoneous plants such as cereals, including wheat and barley.

The amount of combination of the invention to be applied, will depend on various factors such as the compound employed, the subject of the treatment (plant, soil, seed), the type of treatment (e.g. spraying, dusting, seed dressing), the purpose of the treatment (prophylactic or therapeutic), the type of fungi to be treated and the application time.

Particularly preferred mixing partners of the compound I are those which comprise as component b) a compound IIB, IIG, IIH, IIK or IIL.

Another preferred mixing partners of the compound I are those which comprise as component b) a compound IIM, IIN, IIP or IIQ.

Another embodiment of the present invention is represented by those combination which comprise as component a) the compound I and as component b) a compound IIC, IID, IIE, IIF or IIJ.

Another combination is represented by the mixture comprising as component a) the compound I and as component b) the compound of the formula IIA.

It has been found that the use of compound I in combination with the compounds of formula II surprisingly and substantially enhances the effectiveness of the latter against fungi, and vice versa. Additionally, the method of the invention is effective against a wider spectrum of such fungi that can be combated with the active ingredients of this method when used solely.

The weight ratio of a):b) is so selected as to give a synergistic fungicidal action. In general the weight ratio of a) : b) is between 10 : 1 and 1 : 20. The synergistic action of the composition is apparent from the fact that the fungicidal action of the composition of a) + b) is greater than the sum of the fungicidal actions of a) and b).

Where the component b) is the compound IIA the weight ratio of a):b) is for example between 6:1 and 1:6, especially 2:1 and 1:2.

Where the component b) is a compound of formula IIB the weight ratio of a):b) is for example between 5:1 and 1:20, especially 2:1 and 1:20, and more preferably 1:1 to 1:10.

Where component b) is a compound of formula IIC, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is the compound IID, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is the compound IIE, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is a compound of formula IIF, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is a compound of formula IIG, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.



Where component b) is a compound of formula IIH, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is the compound IIJ, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is a compound of formula IIK, the weight ratio of a) : b) is for example between 5:1 and 1:20, especially 3:1 and 1:10, and preferably 2:1 and 1:5.

Where component b) is the compound IIL, the weight ratio of a) : b) is for example between 5:1 and 1:5, specially 2:1 and 1:2, and more preferably 1.5:1 and 1:1.5.

Where component b) is the compound IIM, the weight ratio of a) : b) is for example between 5:1 and 1:5, specially 2:1 and 1:2.

Where component b) is the compound IIN, the weight ratio of a) : b) is for example between 6:1 and 1:6, specially 2:1 and 1:2.

Where component b) is the compound IIP, the weight ratio of a) : b) is for example between 6:1 and 1:6, specially 2:1 and 1:2.

Where component b) is the compound IIQ, the weight ratio of a) : b) is for example between 6:1 and 1:6, specially 2:1 and 1:2.

The method of the invention comprises applying to the treated plants or the locus thereof in admixture or separately, a fungicidally effective aggregate amount of compound I and a compound of component b).

The term locus as used herein is intended to embrace the fields on which the treated crop plants are growing, or where the seeds of cultivated plants are sown, or the place where the seed will be placed into the soil. The term seed is intended to embrace plant propagating material such as cuttings, seedlings, seeds, germinated or soaked seeds.

The novel combinations are extremely effective on a broad spectrum of phytopathogenic fungi, in particular from the Fungi imperfecti and Oomycetes classes. Some of them have a systemic action and can be used as foliar and soil fungicides.

The fungicidal combinations are of particular interest for controlling a large number of fungi in various crops or their seeds, especially wheat, rye, barley, oats, rice, maize, lawns, cotton, soybeans, coffee, sugarcane, fruit and ornamentals in horticulture and viticulture, and in vegetables such as cucumbers, beans and cucurbits.

The combinations are applied by treating the fungi or the seeds, plants or materials threatened by fungus attack, or the soil with a fungicidally effective amount of the active ingredients.

The agents may be applied before or after infection of the materials, plants or seeds by the fungi.

The novel combinations are particularly useful for controlling the following plant diseases:

*Erysiphe graminis* in cereals,

*Erysiphe cichoracearum* and *Sphaerotheca fuliginea* in cucurbits,

*Podosphaera leucotricha* in apples,

*Uncinula necator* in vines,

*Puccinia* species in cereals,

*Rhizoctonia* species in cotton, rice and lawns,

*Ustilago* species in cereals and sugarcane,

*Venturia inaequalis* (scab) in apples,

*Helminthosporium* species in cereals,

*Septoria nodorum* in wheat,

*Septoria tritici* in wheat wheat,

*Rhynchosporium secalis* on barley

*Botrytis cinerea* (gray mold) in strawberries, tomatoes and grapes,

*Cercospora arachidicola* in groundnuts,

*Peronospora tabacina* on tobacco,

*Bremia lactucae* on lettuce,

*Pythium debaryanum* on sugar beet,

*Pseudocercospora herpotrichoides* (*Tapesia* spp.) in wheat and barley,

*Pyrenophora teres* in barley

*Pyricularia oryzae* in rice,

*Phytophthora infestans* in potatoes and tomatoes,

*Fusarium* and *Verticillium* species in various plants,

Plasmopara viticola in grapes,  
Alternaria species in fruit and vegetables.

When applied to the plants the compound I is applied at a rate of 50 to 200 g/ha, particularly 75 to 150 g/ha, e.g. 75, 100, or 125g/ha, in association with 50 to 1500 g/ha, particularly 60 to 1000 g/ha, e.g. 75 g/ha, 80 g/ha, 100 g/ha, 125 g/ha, 150 g/ha, 175 g/ha, 200 g/ha, 300 g/ha, 500 g/ha, or 1000 g/ha of a compound of component b), depending on the class of chemical employed as component b). Where the component b) is the compound IIA for example 50 to 400 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIB for example 50 to 1500 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIC for example 50 to 300 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IID for example 50 to 300 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIE for example 50 to 300 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIF for example 50 to 300 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIG for example 50 to 400 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIH for example 50 to 400 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIJ for example 50 to 300 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIK for example 20 to 2000 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIL for example 50 to 200 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIM for example 50 to 200 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIN for example 50 to 400 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIP for example 50 to 400 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIQ for example 50 to 400 g a.i./ha is applied in association with the compound I.

In agricultural practice the application rates depend on the type of effect desired, and range from 0.02 to 3 kg of active ingredient per hectare.

When the active ingredients are used for treating seed, rates of 0.001 to 50, and preferably from 0.01 to 10g per kg of seed are generally sufficient.

The invention also provides fungicidal compositions comprising the compound I and a compound of component b).

The composition of the invention may be employed in any conventional form, for example in the form of a twin pack, an instant granulate, a flowable or a wettable powder in combination with agriculturally acceptable adjuvants. Such compositions may be produced in conventional manner, e.g. by mixing the active ingredients with appropriate adjuvants (diluent or solvents and optionally other formulating ingredients such as surfactants). Suitable carriers and adjuvants may be solid or liquid and correspond to the substances ordinarily employed in formulation technology, such as, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, thickeners, binding agents or fertilizers. Such carriers are for example described in WO 96/22690.

Particularly formulations to be applied in spraying forms such as water dispersible concentrates or wettable powders may contain surfactants such as wetting and dispersing agents, e.g. the condensation product of formaldehyde with naphthalene sulphonate, an alkylarylsulphonate, a lignin sulphonate, a fatty alkyl sulphate, and ethoxylated alkylphenol and an ethoxylated fatty alcohol.

A seed dressing formulation is applied in a manner known per se to the seeds employing the combination of the invention and a diluent in suitable seed dressing formulation form, e.g. as an aqueous suspension or in a dry powder form having good adherence to the seeds. Such seed dressing formulations are known in the art. Seed dressing formulations may contain the single active ingredients or the combination of active ingredients in encapsulated form, e.g. as slow release capsules or microcapsules.

In general, the formulations include from 0.01 to 90% by weight of active agent, from 0 to 20% agriculturally acceptable surfactant and 10 to 99.99% solid or liquid adjuvant(s), the active agent consisting of at least the compound of formula I together with a compound of component b), and optionally other active agents, particularly guazatin and fenpiclonil. Concentrate forms of compositions generally contain in between about 2 and 80%, preferably between about 5 and 70% by weight of active agent. Application forms of formulation may for example contain from 0.01 to 20% by weight, preferably from 0.01 to 5% by weight of active agent.

Examples for specific formulations-combination are as disclosed e.g. in WO 96/22690, e.g. for wettable powders, emulsifiable concentrate, dusts, extruder granules, coated granules, suspension concentrate.

#### Slow Release Capsule Suspension

28 parts of a combination of the compound I and a compound of component b), or of each of these compounds separately, are mixed with 2 parts of an aromatic solvent and 7 parts of toluene diisocyanate/polymethylene-polyphenylisocyanate-mixture (8:1). This mixture is emulsified in a mixture of 1.2 parts of polyvinylalcohol, 0.05 parts of a defoamer and 51.6 parts of water until the desired particle size is achieved. To this emulsion a mixture of 2.8 parts 1,6-diaminohexane in 5.3 parts of water is added. The mixture is agitated until the polymerization reaction is completed.

The obtained capsule suspension is stabilized by adding 0.25 parts of a thickener and 3 parts of a dispersing agent. The capsule suspension formulation contains 28% of the active ingredients. The medium capsule diameter is 8-15 microns.

The resulting formulation is applied to seeds as an aqueous suspension in an apparatus suitable for that purpose.

Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations.

#### Biological Examples

A synergistic effect exists whenever the action of an active ingredient combination is greater than the sum of the actions of the individual components.

The action to be expected E for a given active ingredient combination obeys the so-called COLBY formula and can be calculated as follows (COLBY, S.R. "Calculating synergistic and antagonistic responses of herbicide combination". Weeds, Vol. 15, pages 20-22; 1967):

ppm = milligrams of active ingredient (= a.i.) per litre of spray mixture

X = % action by active ingredient I using p ppm of active ingredient

Y = % action by active ingredient II using q ppm of active ingredient.

According to Colby, the expected (additive) action of active ingredients I+II using p+q ppm

of active ingredient is  $E = X + Y - \frac{X \cdot Y}{100}$

If the action actually observed (O) is greater than the expected action (E), then the action of the combination is superadditive, i.e. there is a synergistic effect.

Alternatively the synergistic action may also be determined from the dose response curves according to the so-called WADLEY method. With this method the efficacy of the a.i. is determined by comparing the degree of fungal attack on treated plants with that on untreated, similarly inoculated and incubated check plants. Each a.i. is tested at 4 to 5 concentrations. The dose response curves are used to establish the EC90 (i.e. concentration of a.i. providing 90% disease control) of the single compounds as well as of the combinations (EC 90<sub>observed</sub>). The thus experimentally found values of the mixtures at a given weight ratio are compared with the values that would have been found were only a complementary efficacy of the components was present (EC 90 (A+B)<sub>expected</sub>). The EC90 (A+B)<sub>expected</sub> is calculated according to Wadley (Levi et al., EPPO- Bulletin 16, 1986, 651-657):

$$\text{EC 90 (A+B)}_{\text{expected}} = \frac{a + b}{\frac{a}{\text{EC90 (A)}_{\text{observed}}} + \frac{b}{\text{EC90 (B)}_{\text{observed}}}}$$

wherein a and b are the weight ratios of the compounds A and B in the mixture and the indexes (A), (B), (A+B) refer to the observed EC 90 values of the compounds A, B or the given combination A+B thereof. The ratio EC90 (A+B)<sub>expected</sub> / EC90 (A+B)<sub>observed</sub> expresses the factor of interaction (F). In case of synergism, F is >1.

**Example B-1: Residual-protective action against *Venturia inaequalis* on apples**

Apple cuttings with 10-20 cm long fresh shoots are sprayed to drip point with an aqueous spray mixture prepared from a wettable powder formulation of the active ingredient mixture and infected 24 hours later with a conidia suspension of the fungus. The plants are incubated for 5 days at 90-100 % relative humidity and stood in a greenhouse for a further 10 days at 20-24°C. Fungus infestation is evaluated 12 days after infection.

**Example B-2(a): Action against *Botrytis cinerea* on apple fruits**

Artificially damaged apples are treated by dropping a spray mixture of the active ingredient mixture onto the damage sites. The treated fruits are then inoculated with a spore suspension of the fungus and incubated for one week at high humidity and about 20°C. The fungicidal action of the test compound is derived from the number of damage sites that have begun to rot.

Example B-2(b): Action against *Botritis cinerea* on tomatoes

4 week old tomato plants cv. "Roter Gnom" were treated with the formulated test compound in a spray chamber. Two days after application the tomato plants were inoculated by spraying a spore suspension on the test plants. After an incubation period of 4 days at 20°C and 95% relative humidity in a growth chamber the disease incidence was assessed.

Example B-2(c): Action against *Botritis cinerea* on grapes

5 week old grape seedlings cv. "Gutedel" were treated with the formulated test compound in a spray chamber. Two days after application the grape plants were inoculated by spraying a spore suspension on the test plants. After an incubation period of 4 days at 21°C and 95% relative humidity in a greenhouse the disease incidence was assessed.

Example B-3: Action against *Podosphaera leucotricha* on apple shoots

Apple cuttings with about 15 cm long fresh shoots are sprayed with a spray mixture of the active ingredient mixture. The treated plants are infected 24 hours later with a conidia suspension of the fungus and placed in a climatic chamber at 70 % relative humidity and 20°C. Fungus infestation is evaluated 12 days after infection.

Example B-4: Action against *Drechslera teres* on barley

10-day-old barley plants of the "Golden Promise" variety are sprayed with a spray mixture of the active ingredient mixture. The treated plants are infected 24 hours later with a conidia suspension of the fungus and incubated in a climatic chamber at 70 % relative humidity and 20-22°C. Fungus infestation is evaluated 5 days after infection.

Example B-5 : Efficacy against *Erysiphe graminis* f.sp. *tritici* on wheat

Five to ten wheat seeds c.v. "Arina" are sown in plastic pots of 7 cm diameter and grown for 7 to 12 days at 20°C, 50-70% rH. When the primary leaves have fully expanded, the plants

are spray treated with aqueous spray liquors containing the single compounds, or mixtures thereof (hereinafter a.i.). All compounds are used as experimental or commercially available formulations, combinations are applied as tank mixtures. The application comprises foliar spraying to near runoff (three pots per treatment). 24 hours after the application or 24 hours before application, the plants are inoculated in a settling tower with fresh spores of *Erysiphe graminis* f. sp. *tritici*. The plants are then incubated in a growth chamber at 20°C, 60% rH. Seven days after the inoculation, the percentage of infection on primary leaves is evaluated. The efficacy of the a.i. is determined by comparing the degree of fungal attack on treated plants with that on untreated, similarly inoculated and incubated check plants. Each a.i. is tested at 4 to 5 concentrations. The synergy factor is calculated according to the COLBY method.

Example B-7: Activity against Uncinula necator

Grape plants, 4 weeks old (4-5 leaves), are sprayed to near run off with a suspension containing 250 mg/l of active ingredient. The deposit is then allowed to dry. One day later, the treated plants are inoculated by dusting freshly harvested conidia over the test plants; then the plants were incubated in a growth chamber for 10-14 days at +22°C and 70% r.h. The efficacy of the test compounds is determined by comparing the degree of fungal attack with that on untreated, similarly inoculated check plants.

The mixtures according to the invention exhibit good activity in these Examples.

Example B-8: Activity against Plasmopara viticola in grapevines

Grapevine seedlings at the 4- to 5-leaf stage are sprayed to drip point with an aqueous spray mixture prepared with a wettable powder of the active ingredient mixture (0.02% of active ingredient) and, 24 hours later, infected with a sporangia suspension of the fungus. The fungus infestation is assessed 6 days after infection, during which time a relative atmospheric humidity of 95 to 100% and a temperature of 20°C are maintained.

Example B-9: Activity against Phytophthora infestans in tomatoes

a) Curative action

Tomato plants cv. "Roter Gnom" are grown for three weeks and then sprayed with a zoospore suspension of the fungus and incubated in a cabin at 18 to 20°C and saturated atmospheric humidity. The humidification is interrupted after 24 hours. After the plants have dried, they are sprayed with a mixture which comprises the active ingredient formulated as a



wettable powder at a concentration of 200ppm. After the spray coating has dried, the plants are returned to the humid chamber for 4 days. Number and size of the typical foliar lesions which have appeared after this time are used as a scale for assessing the efficacy of the test substances.

b) Preventive-systemic action

The active ingredient which is formulated as a wettable powder is introduced, at a concentration of 60ppm (relative to the soil volume), onto the soil surface of three-week-old tomato plants cv. "Roter Gnom" in pots. After an interval of three days, the underside of the leaves is sprayed with a zoospore suspension of *Phytophthora infestans*. They are then kept for 5 days in a spray cabin at 18 to 20°C and saturated atmospheric humidity. After this time, typical foliar lesions appear whose number and size are used for assessing the efficacy of the test substances.

Example B-10: Activity against *Phytophthora* in potato plants

a) Residual-protective action

2-3 week old potato plants (Bintje variety) are grown for 3 weeks and then sprayed with a spray mixture (0.02% of active ingredient) prepared with a wettable powder of the active ingredient. After 24 hours, the treated plants are infected with a sporangia suspension of the fungus. The fungus infestation is assessed after the infected plants have been incubated for 5 days at a relative atmospheric humidity of 90-100% and 20°C.

b) Systemic action

A spray mixture (0.002% of active ingredient based on the soil volume) prepared with a wettable powder of the active ingredient is poured next to 2-3 week old potato plants (Bintje variety) which have been grown for 3 weeks. Care is taken that the spray mixture does not come into contact with the aerial parts of the plants. After 48 hours, the treated plants are infected with a sporangia suspension of the fungus. Fungus infestation is assessed after the infected plants have been incubated for 5 days at a relative atmospheric humidity of 90-100% and 20°C.

The efficacy of the test combinations and the single active ingredients in the above tests is determined by comparing the degree of fungal attack with that on untreated, similarly inoculated check plants.

The mixtures according to the invention exhibit good activity in these Examples.


## WHAT IS CLAIMED IS:

1. A method of combating phytopathogenic diseases on crop plants which comprises applying to the crop plants or the locus thereof being infested with said phytopathogenic disease, in admixture or separately, an effective amount of a combination of a first component


- a) 2-[ $\alpha$ -{[( $\alpha$ -methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acid methyl ester-O-methyloxime (I)

in association with

- b) a second component selected from either  
a compound of formula IIA




an anilide of formula IIB



wherein R<sub>1</sub> is fluorine or chlorine; or

a carbamate of formula IIC



wherein X is N or CH, and R<sub>2</sub> is 4-CH<sub>3</sub>, 4-Cl or 2,4-dichloro; or


a compound IID

methyl(2)-2-[2-[6-(trifluoromethyl)pyrid-2-yloxymethyl]-phenyl]-3-methoxyacrylate; or

a compound IIE


(E)-N-methyl-2-[2-(2,5-dimethylphenoxy-methyl)phenyl]-2-methoxy-iminoacetamide; or

a compound of formula IIF



wherein R<sub>3</sub> is methyl or ethyl; or

a (S)-valinamide of formula IIG




wherein R<sub>4</sub> is isopropyl, sec.-butyl or tert.-butyl, and

R<sub>5</sub> is 4-chlorophenyl, 4-methylphenyl, 4-methoxyphenyl or β-naphthyl, and

wherein the asymmetric center is preferably (R); or

a (S)-valinamide of formula IIH




wherein R<sub>6</sub> is isopropyl, sec.-butyl or tert.-butyl, R<sub>7</sub> is halogen, methyl or methoxy and

n is 0, 1 or 2; or

a compound IIJ

N-methyl-2-[2-( $\alpha$ -methyl-3-(trifluoromethyl)benzyloximinomethyl)phenyl]-2-methoximinoacetamide; or

a compound of formula IIK



wherein  $R_8$  is halogen or  $C_1$ - $C_4$ -alkyl, preferably chlorine; or


a compound IIL

N-(3'-(1'-chloro-3-methyl 2'-oxopentane))-3,5-dichloro-4-methylbenzamide; or


a compound IIM

(S)-1-anilino-4-methyl-2-methylthio-4-phenylimidazolin-5-one; or

a compound of formula IIN



a compound of formula IIP



a compound IIQ

N-methyl-2-[ $\alpha$ -{[( $\alpha$ -methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acidamide-O-methyloxime.

2. A method according to claim 1 wherein the components b) is selected from the compounds of formula IIB, IIG, IIH, IIK and IIL.
3. A method according to claim 1 wherein the component b) is selected from the group comprising IIC, IID, IIE, IIF and IIJ.
4. A method according to claim 1 wherein the component b) is a compound of formula IIA.
5. A method according to claim 1 wherein the component b) is selected from the group comprising IIM, IIN, IIP or IIQ.
6. A method according to any of claims 1 to 5 wherein the components a) and b) are applied in a quantity producing a synergistic disease controlling effect, specially a fungicidal effect.
7. A fungicidal composition comprising a fungicidally effective combination of
  - a) a compound I according to claim 1  
in association with
  - b) either a compound IIA, or  
a compound of formula IIB, or  
a compound of formula IIC, or  
a compound IID, or  
a compound IIE, or  
a compound of formula IIF, or  
a compound of formula IIG, or  
a compound of formula IIH, or  
a compound IIJ, or  
a compound of formula IIK, or  
a compound IIL, or

a compound of formula IIM, or  
a compound of formula IIN, or  
a compound of formula IIP, or  
a compound IIQ  
as defined in claim 1.

8. A composition according to claim 7 the weight ratio of a) to b) is between 10 : 1 and 1 : 20.

9. A composition according to claim 7 wherein the component b) is selected from the group comprising IIB, IIG, IIH, IIK and IIL.

10. A composition according to claim 7 wherein the component b) is selected from the group comprising IIC, IID, IIE, IIF and IIJ.

11. A composition according to claim 7 wherein the component b) is a compound of formula IIA.

12. A composition according to claim 7 wherein the component b) is selected from the group comprising IIM, IIN, IIP and IIQ.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> A01N 37/50 // (A01N 37/50, 47:24, 47:12, 43:88, 43:653, 43:50, 43:40, 37:52, 37:46, 37:24, 37:20)		<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/63813</b>  <b>(43) International Publication Date:</b> 16 December 1999 (16.12.99)
<b>(21) International Application Number:</b> PCT/EP99/03883  <b>(22) International Filing Date:</b> 4 June 1999 (04.06.99)  <b>(30) Priority Data:</b> 9812331.8      8 June 1998 (08.06.98)      GB 9903669.1      17 February 1999 (17.02.99)      GB  <b>(71) Applicant (for all designated States except AT US):</b> NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).  <b>(71) Applicant (for AT only):</b> NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> ZURFLÜH, René [CH/CH]; Dachslenbergstrasse 54, CH-8180 Bülach (CH). LEADBITTER, Neil [GB/CH]; Therwilerstrasse 117, CH-4104 Oberwil (CH).  <b>(74) Agent:</b> BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> 2 March 2000 (02.03.00)	
<b>(54) Title:</b> FUNGICIDAL COMBINATIONS COMPRISING A GLYOXALIC ACID METHYLESTER METHYLOXIME DERIVATIVE			
<b>(57) Abstract</b>  A method of combating phytopathogenic diseases on crop plants which comprises applying to the crop plants or the locus thereof being infested with said phytopathogenic disease an effective amount of a combination of a) 2-[ $\alpha$ -{[( $\alpha$ -methyl-3-trifluoromethyl-benzyl)imino]-oxy}- $\alpha$ -tolyl]-glyoxalic acid methyl ester-O-methyloxime (I) in association with b) a broad variety of other plant fungicides is particularly effective in combating or preventing diseases of crop plants. These combinations exhibit synergistic fungicidal activity.			

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		



# INTERNATIONAL SEARCH REPORT

Internat I Application No

PCT/EP 99/03883

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01N37/50

43:88,43:653,43:50,43:40,37:52,

/(A01N37/50,47:24.47:12,  
37:46.37:24,37:20)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97 40671 A (CIBA GEIGY AG ; ZEUN RONALD (DE)) 6 November 1997 (1997-11-06) page 1 -page 2, paragraph 2; claims 1,12; examples B-1,B-4,B-6,B-8,B-11 ---	1,4,6-8, 11
Y	DE 195 43 746 A (BASF AG) 28 May 1997 (1997-05-28) page 1 -page 3 page 4, table I.2, compound no. I.2.7 page 6, line 32 - line 37 page 9, table II.2C, compound no. II.2C-4 page 16, line 64 -page 21, line 60; claims --- -/-	1,4,6-8, 11

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

5 October 1999

Date of making of the international search report

11. 01. 2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

MÜLLNERS W.

# INTERNATIONAL SEARCH REPORT

Intern: al Application No

PCT/EP 99/03883

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation or document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 00011 A (CIBA GEIGY AG ;KNAUF BEITER GERTRUDE (DE); ZEUN RONALD (DE)) 3 January 1997 (1997-01-03) page 1 -page 3, line 2; claims 1,13; examples B-1,B-2,B-4,B-5,B-8,B-11,B-12,B-13 ---	1,4,6-8, 11
A	WO 97 00012 A (CIBA GEIGY AG ;KNAUF BEITER GERTRUDE (DE); KUENG RUTH BEATRICE (CH)) 3 January 1997 (1997-01-03) page 1 -page 2, paragraph 1 page 3, paragraph 2; claims 1,13; examples B-1,B-4,B-6,B-8,B-11 ---	1,4,6-8, 11
A	EP 0 626 135 A (BAYER AG) 30 November 1994 (1994-11-30) page 2, line 1 - line 30 page 7, line 25 - line 45; claims; examples 3-7 ---	1,4,6-8, 11
A	DE 44 37 048 A (BAYER AG) 18 April 1996 (1996-04-18) page 2, line 1 - line 23 page 1, line 46 - line 47 page 10 -page 11; claim 1 ---	1,4,6-8, 11
P,X	WO 98 25459 A (CIBA GEIGY AG ;MARGOT PAUL (CH); KNAUF BEITER GERTRUDE (DE)) 18 June 1998 (1998-06-18) page 1 -page 2, paragraph 1; claim 1 -----	1,4,6-8, 11

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP 99/03883

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
  
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1, 6-8 (partially); 4, 11 (completely)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## 1. Claims: 1, 6-8 (partially); 4, 11 (completely)

A method of combating phytopathogenic diseases on crop plants which comprises applying to the plants or the locus thereof, in admixture or separately, a combination of a first component (A) which is  
2-[alpha-[[alpha-methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acid methylester methyloxime  
in association with a second component (B) which is  
the cyclohexancarboxanilide of formula (IIA)  
and corresponding fungicidal compositions

## 2. Claims: 1, 2, 6-9 (all partially)

As subject 1 but the second component (B) is  
the N-biphenyl-2-chlor-nicotinamide of formula (IIB)

## 3. Claims: 1, 5-8, 12 (partially); 3, 10 (completely)

As subject 1 but the second component (B) is  
a strobilurin fungicide selected from the list consisting of  
the compounds IIC, IID, IIE, IIF, IIJ, IIN, IIP and IIQ

## 4. Claims: 1, 2, 6-9 (all partially)

As subject 1 but the second component (B) is  
a valinamide fungicide selected from the list consisting of  
the compounds IIG and IIH

## 5. Claims: 1, 2, 6-9 (all partially)

As subject 1 but the second component (B) is  
a 4-(2-Fluoroalkylamido)-benzamide of formula (IIK)

## 6. Claims: 1, 2, 6-9 (all partially)

As subject 1 but the second component (B) is  
the N-acetonyl-benzamide (IIL)

## 7. Claims: 1, 5-8, 12 (all partially)

As subject 1 but the second component (B) is  
the phenylimidazolinone (IIM)

## INTERNATIONAL SEARCH REPORT

Internat. Application No.

PCT/EP 99/03883

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9740671 A	06-11-1997	AU 2696697 A	19-11-1997
DE 19543746 A	28-05-1997	AU 7626896 A	19-06-1997
		BR 9611631 A	25-05-1999
		CA 2235039 A	05-06-1997
		CN 1202798 A	23-12-1998
		CZ 9801610 A	14-10-1998
		WO 9719595 A	05-06-1997
		EP 0863702 A	16-09-1998
		NZ 322502 A	28-01-1999
		SK 50798 A	02-12-1998
WO 9700011 A	03-01-1997	AU 709775 B	09-09-1999
		AU 6300096 A	15-01-1997
		BR 9608356 A	18-08-1998
		CA 2221759 A	03-01-1997
		CZ 9704041 A	13-05-1998
		EP 0831697 A	01-04-1998
		HU 9801819 A	30-11-1998
		JP 11507654 T	06-07-1999
		PL 323677 A	14-04-1998
		ZA 9605081 A	17-12-1996
WO 9700012 A	03-01-1997	AU 690464 B	23-04-1998
		AU 6125196 A	15-01-1997
		BR 9608358 A	18-08-1998
		CA 2224977 A	03-01-1997
		CZ 9704040 A	13-05-1998
		EP 0831698 A	01-04-1998
		HU 9802198 A	28-01-1999
		JP 11507655 T	06-07-1999
		NZ 310452 A	29-09-1999
		PL 323945 A	27-04-1998
		ZA 9605080 A	17-12-1996
EP 0626135 A	30-11-1994	DE 4313867 A	03-11-1994
		CN 1094221 A	02-11-1994
		DE 59402194 D	30-04-1997
		ES 2099501 T	16-05-1997
		JP 6329505 A	29-11-1994
		TR 27417 A	14-04-1995
		US 5532262 A	02-07-1996
		US 5672619 A	30-09-1997
		US 5922762 A	13-07-1999
		ZA 9402887 A	04-01-1995
DE 4437048 A	18-04-1996	AT 175839 T	15-02-1999
		AU 699963 B	17-12-1998
		AU 3317395 A	02-05-1996
		BR 9504413 A	27-05-1997
		CN 1132591 A	09-10-1996
		DE 59504892 D	04-03-1999
		EP 0707792 A	24-04-1996
		ES 2129168 T	01-06-1999
		GR 3029400 T	28-05-1999
		HU 73550 A	28-08-1996
		JP 8208406 A	13-08-1996
		PL 310937 A	29-04-1996

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/03883

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 4437048 A		US 5962518 A	05-10-1999
		ZA 9508711 A	22-05-1996
-----			
WO 9825459 A	18-06-1998	AU 5756398 A	03-07-1998
		CZ 9902122 A	15-09-1999
		EP 0944308 A	29-09-1999
		NO 992750 A	07-06-1999
-----			

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> A01N 37/50 // (A01N 37/50, 47:24, 47:12, 43:88, 43:653, 43:50, 43:40, 37:52, 37:46, 37:24, 37:20)		<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/63813</b> <b>(43) International Publication Date:</b> 16 December 1999 (16.12.99)
<b>(21) International Application Number:</b> PCT/EP99/03883 <b>(22) International Filing Date:</b> 4 June 1999 (04.06.99) <b>(30) Priority Data:</b> 9812331.8 8 June 1998 (08.06.98) GB 9903669.1 17 February 1999 (17.02.99) GB <b>(71) Applicant (for all designated States except AT US):</b> NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). <b>(71) Applicant (for AT only):</b> NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> ZURFLÜH, René [CH/CH]; Dachslenbergstrasse 54, CH-8180 Bülach (CH). LEADBITTER, Neil [GB/CH]; Therwilerstrasse 117, CH-4104 Oberwil (CH). <b>(74) Agent:</b> BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).			<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>With amended claims.</i>  <b>(88) Date of publication of the international search report:</b> 2 March 2000 (02.03.00)  <b>Date of publication of the amended claims:</b> 13 April 2000 (13.04.00)
<b>(54) Title:</b> FUNGICIDAL COMBINATIONS COMPRISING A GLYOXALIC ACID METHYLESTER METHYLOXIME DERIVATIVE			
<b>(57) Abstract</b> <p>A method of combating phytopathogenic diseases on crop plants which comprises applying to the crop plants or the locus thereof being infested with said phytopathogenic disease an effective amount of a combination of a) 2-[α-{{[(α-methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acid methyl ester-O-methyloxime (I) in association with b) a broad variety of other plant fungicides is particularly effective in combating or preventing diseases of crop plants. These combinations exhibit synergistic fungicidal activity.</p>			

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						



**AMENDED CLAIMS**


[received by the International Bureau on 25 February 2000 (25.02.00);  
original claims 1 and 7 amended; original claims 4 and 11 cancelled;  
remaining claims unchanged (4 pages)]

1. A method of combating phytopathogenic diseases on crop plants which comprises applying to the crop plants or the locus thereof being infested with said phytopathogenic disease, in admixture or separately, an effective amount of a combination of a first component

a) 2-[ $\alpha$ -{[( $\alpha$ -methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acid methyl ester-O-methyloxime (I)


in association with

b) a second component selected from either  
an anilide of formula IIB



wherein  $R_1$  is fluorine or chlorine; or

a carbamate of formula IIC



wherein X is N or CH, and  $R_2$  is 4-CH<sub>3</sub>, 4-Cl or 2,4-dichloro; or


a compound IID

methyl(2)-2-{2-[6-(trifluoromethyl)pyrid-2-yloxymethyl]-phenyl}-3-methoxyacrylate; or

a compound IIE


(E)-N-methyl-2-[2-(2,5-dimethylphenoxy)methyl]phenyl]-2-methoxy-iminoacetamide; or

a compound of formula IIF



wherein R<sub>3</sub> is methyl or ethyl; or

a (S)-valinamide of formula IIG




wherein R<sub>4</sub> is isopropyl, sec.-butyl or tert.-butyl, and

R<sub>5</sub> is 4-chlorophenyl, 4-methylphenyl, 4-methoxyphenyl or β-naphthyl, and

wherein the asymmetric center is preferably (R); or

a (S)-valinamide of formula IIH




wherein R<sub>6</sub> is isopropyl, sec.-butyl or tert.-butyl, R<sub>7</sub> is halogen, methyl or methoxy and n is 0,1 or 2; or

a compound IIJ

N-methyl-2-[2-{α-methyl-3-(trifluoromethyl)benzyloximinomethyl}phenyl]-2-methoximinoacetamide; or

a compound of formula IIK



wherein R<sub>8</sub> is halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl, preferably chlorine; or


a compound IIL

N-(3'-(1'-chloro-3-methyl 2'-oxopentan))-3,5-dichloro-4-methylbenzamide; or


a compound IIM

(S)-1-anilino-4-methyl-2-methylthio-4-phenylimidazolin-5-one; or

a compound of formula IIN



a compound of formula IIP



a compound IIQ

N-methyl-2-[α-[[[(α-methyl-3-trifluoromethyl-benzyl)imino]-oxy]-o-tolyl]-glyoxalic acidamide-O-methyloxime.

2. A method according to claim 1 wherein the components b) is selected from the compounds of formula IIB, IIG, IIH, IIK and IIL.
3. A method according to claim 1 wherein the component b) is selected from the group comprising IIC, IID, IIE, IIF and IIJ.
5. A method according to claim 1 wherein the component b) is selected from the group comprising IIM, IIN, IIP or IIQ.
6. A method according to any of claims 1 to 5 wherein the components a) and b) are applied in a quantity producing a synergistic disease controlling effect, specially a fungicidal effect.
7. A fungicidal composition comprising a fungicidally effective combination of
  - a) a compound I according to claim 1  
in association with
  - b) either a compound of formula IIB, or  
a compound of formula IIC, or  
a compound IID, or  
a compound IIE, or  
a compound of formula IIF, or  
a compound of formula IIG, or  
a compound of formula IIH, or  
a compound IIJ, or  
a compound of formula IIK, or  
a compound IIL, or  
a compound of formula IIM, or  
a compound of formula IIN, or  
a compound of formula IIP, or  
a compound IIQ  
as defined in claim 1.
8. A composition according to claim 7 the weight ratio of a) to b) is between 10 : 1 and 1 : 20.